CHAPTER - 1

NUMBER SYSTEMS

- 1. Rational Numbers
- 2. Irrational Numbers
- 3. Real Numbers and their Decimal Expansions
- 4. Operations on Real Numbers
- 5. Laws of Exponents for Real Numbers



- Natural numbers are 1, 2, 3, denoted by N.
- Whole numbers are 0, 1, 2, 3, denoted by W.
- Integers -3, -2, -1, 0, 1, 2, 3, denoted by Z.
- Rational numbers All the numbers which can be written in the form r/s p/q, are called rational numbers where p and q are integers.
- Irrational numbers A number s is called irrational, if it cannot be written in the form p/q where p and q are integers and
- The decimal expansion of a rational number is either terminating or non-terminating recurring. Thus we say that a number whose decimal expansion is either terminating or non-terminating recurring is a rational number.
- The decimal expansion of a irrational number is non terminating non-recurring.
- All the rational numbers and irrational numbers taken together.
- Make a collection of real number.
- A real no is either rational or irrational.
- If r is rational and s is irrational then r+s, r-s, r.s are always irrational numbers but r/s may be rational or irrational.
- Every irrational number can be represented on a number line using Pythagoras theorem.
- Rationalization means to remove square root from the denominator.

 $\frac{3+\sqrt{5}}{\sqrt{2}}$ to remove we will multiply both numerator & denominator by $\sqrt{2}$ $\frac{1}{a\pm\sqrt{b}}$ its rationalization factor $a\mp\sqrt{b}$